B P International

ABOUT

Search

Theory and Applications of Mathematical Science Vol. 2

Dr. Narayan Thapa, (Editor)
Associate Professor, Department of Mathematical Sciences, Cameron University, Lawton, Oklahoma, USA.

Profile Link: http://www.cameron.edu/math/faculty.html

Short Biosketch

Charles Roberto Telles
Department of Research Advisory, Secretary of State for Education and Sport, Curitiba, Paraná, Brazil.

K. Mahesh
Department of Electrical and Electronics Engineering, Sir M Visvesvaraya Institute of Technology, Bengaluru, India.

J. Lithesh
Department of Electrical and Electronics Engineering, New Horizon College of Engineering, Bengaluru, India.

Sukumar Mondal
Department of Mathematics, Raja N. L. Khan Women's College (Autonomous), Gope Palace, Paschim Medinipur, 721 102, West Bengal, India.

Mohammed Barkatou
Innovation in Sciences, Technology and Modeling Laboratory, University of Chouaïb Doukkali, Morocco.

Saif Ur Rehman
Department of Mathematics, Gomal University, Dera Ismail Khan 29050, Pakistan.

Mohammed Harfaoui
University Hassan II Mohammedia, Laboratory of Mathematics, Criptography and Mechanical F. S. T., BP 146, Mohammedia 20650, Morocco.

Loubna Lakhmaili
University Hassan II Mohammedia, Laboratory of Mathematics, Criptography and Mechanical F. S. T., BP 146, Mohammedia 20650, Morocco.

Abdellah Mourassil
University Hassan II Mohammedia, Laboratory of Mathematics, Criptography and Mechanical F. S. T., BP 146, Mohammedia 20650, Morocco.

L. I. Petrova
Moscow State University, Russia.

Kavita Sakure
Department of Mathematics, Govt. Digvijay Auto. P.G. College, Rajnandgaon, 491441, India.

Juhani Riihentaus
Department of Mathematical Sciences, University of Oulu, P.O. Box 3000, FI-90014 Oulun Yliopisto, Finland and Department of Physics and Mathematics, University of Eastern Finland, P.O. Box 111, FI-80101 Joensuu, Finland.

Ahmad Alghoussein
Department of Mathematics, Faculty of Science, Tishreen University, Lattakia, Syria.

Ziad Kanaya
Department of Mathematics, Faculty of Science, Tishreen University, Lattakia, Syria.

Salwa Yacoub
Department of Mathematics, Faculty of Science, Tishreen University, Lattakia, Syria.

Synopsis

This book covers all areas of mathematical science. The contributions by the authors include nonlinear integral equation; Darbo's fixed point theorem; Weibull parameters; MATLAB; trapezoid graphs; fixed point; common fixed point; fuzzy cone metric space; Dirichlet problem; Quadrature surfaces; agent-based modelling; nonlinear dynamics; topological groups; free topological groups; nonconsistency of the conservation laws equations; degenerate transformation; discrete solutions; meromorphic functions; Pad-approximants; (p, q)-order and (p, q)-type; logarithmic capacity; quasinearly subharmonic; families of quasinearly subharmonic functions etc. This book contains various materials suitable for students, researchers and academicians in the field of mathematical science.



Media Promotion: Chapter 01

Media Promotion: Chapter 02

Media Promotion: Chapter 03

Media Promotion: Chapter 04

Media Promotion: Chapter 05

Media Promotion: Chapter 06

Media Promotion: Chapter 07

Media Promotion: Chapter 08

Media Promotion: Chapter 09

Media Promotion: Chapter 10

DOI: https://doi.org/10.9734/bpi/tams/v2

ISBN-13 (15)
978-93-89816-39-6

Article Metrics

PLUMX

See Details

BROWSE

New Releases

B P International



Evolutionary Computing and Mobile Sustainable Networks pp 103–112

Review of Python for Solar Photovoltaic Systems

R. Sivapriyan, D. Elangovan & Kavyashri S. N. Lekhana

Conference paper | First Online: 01 August 2020

Part of the Lecture Notes on Data Engineering and Communications Technologies book series (LNDECT,volume 53)

Abstract

In recent years, the usage of solar energy as a source to produce power has increased exponentially as it provides a clean and efficient alternative to depleting non-renewable resources. The normal working period of a photovoltaic (PV) panel is 20 years, but due to defects in manufacturing or atmospheric condition changes, the efficiency and the lifespan of the panel decrease each year. The objective of this review article is to present and analyze the different methods that can be used to reduce the degradation rate of the PV cells in an economically viable way. Open-source frameworks are important to make any solution affordable; hence we explore the usage of python language in developments relating to improvement in the performance of PV cells. Based on this review a practically employable solution to improve working conditions for PV cells can be obtained.

Keywords

Solar energy **PV panels** **Python**

Modeling **Monitoring** **Analysis**

Fault detection

References

1. Fernandes TR, Fernandes LR, Ricciardi TR, Ugarte LF, de Almeida MC (2018) Python Programming Language for Power System Analysis Education and Research. In: IEEE PES Transmission & Distribution Conference and Exhibition-Latin America (T&D-LA), pp. 1–5. IEEE

2. Grout I (2018) Electronic Circuit and System Design using Python and VHDL. In: 15th International Conference on Electrical Engineering/Electronics, Computer, Telecommunications and Information Technology, pp. 13–16. IEEE

3. Bonneel P, Le Besnerais J, Pile R, Devillers E (2018) Pyleecan: an open-source Python object-oriented software for the multiphysic design optimization

of electrical machines. In: XIII International Conference on Electrical Machines, pp 948–954. IEEE

4. Umahsnakar S, Arunkumar G, Sivapriyan R (2017) Comparative analysis of solar photovoltaic cell models. In: 2017 International Conference On Smart Technologies for Smart Nation, pp 46–51. IEEE

5. Sivapriyan R, S. Umashankar, P. Sanjeevikumar, and Atif Iqbal.: Direct-Coupled Permanent Magnet DC Motor-Driven Solar Photovoltaic Water Pumping System—A Literature Review. In: Advances in Smart Grid and Renewable Energy, pp. 307–314. Springer, Singapore (2018)

6. White, Jeremy T., Michael N. Fienen, and John E. Doherty.: A python framework for environmental model uncertainty analysis. In: Environmental Modelling & Software 85. pp. 217–228. (2016)

7. Alonso-Álvarez D, Wilson T, Pearce P, Führer M, Farrell D, Ekins-Daukes N (2018) Solcore: a multi-scale, Python-based library for modelling solar cells and semiconductor materials. J Comput Electron 17(3):1099–1123

8. Charles, Nathan, Mahmoud Kabalan, and Pritpal Singh.: Open source photovoltaic system performance modeling with python. In: IEEE Canada International Humanitarian Technology Conference, pp. 1–4. IEEE (2015)

9. Stein, Joshua S., William F. Holmgren, Jessica Forbess, and Clifford W. Hansen.: PVLIB: Open source photovoltaic performance modeling functions for Matlab and Python. In: ieee 43rd photovoltaic specialists conference, pp. 3425–3430. IEEE (2016)

10. Thurner, Leon, Alexander Scheidler, Florian Schäfer, Jan-Hendrik Menke, Julian Dollichon, Friederike Meier, Steffen Meinecke, and Martin Braun.: Pandapower—An open-source python tool for convenient modeling, analysis, and optimization of electric power systems. In: Transactions on Power Systems 33, no. 6, pp. 6510–6521 (2018)

11. Phongtrakul, Tipthacha, Yuttana Kongjeen, and Krischonme Bhumkittipich.: Analysis of Power Load Flow for Power Distribution System based on PyPSA Toolbox. In: 15th International Conference on Electrical Engineering/Electronics,

Computer, Telecommunications and Information Technology, pp. 764–767. IEEE (2018)

12. Kang, Shinyuk, and Ilwoo Lee.: Implementation of PV Monitoring System Using Python. In: 21st International Conference on Advanced Communication Technology, pp. 453–455. IEEE, (2019)

13. Deshmukh, Neha S., and D. L. Bhuyar.: A Smart Solar Photovoltaic Remote Monitoring and Controlling. In: Second International Conference on Intelligent Computing and Control Systems, pp. 67–71. IEEE (2018)

14. Walters, Joseph, Siyu Guo, Eric Schneller, Hubert Seigneur, and Matthew Boyd.: PV module loss analysis using system in-situ monitoring data. In: IEEE 7th World Conference on Photovoltaic Energy Conversion, pp. 2204–2208. IEEE (2018)

15. Prasanna, J. Laxmi, D. Lavanya, and T. Anil Kumar.: Condition monitoring of a virtual solar system using IoT. In: 2nd International Conference on Communication and Electronics Systems, pp. 286–290. IEEE (2017)

16. Dyamond WP, Rix AJ (2019) Detecting Anomalous Events for a Grid Connected PV Power Plant Using Sensor Data. In: Southern African Universities Power Engineering Conference/Robotics and Mechatronics/Pattern Recognition Association of South Africa, pp. 287–292. IEEE

Acknowledgements

The authors would like to thank Punitharaj Research and Training Institute (OPC) Private Limited, Bangalore (https://prti.co.in) and Sir MVIT, Bangalore for their technical and financial support for this work.

Author information

Authors and Affiliations

Sir MVIT, Bengaluru, India

R. Sivapriyan & Kavyashri S. N. Lekhana

SELECT, VIT, Vellore, India

D. Elangovan

Corresponding author

Correspondence to R. Sivapriyan .

Editor information

Editors and Affiliations

Research and Industry Incubation Center, Department of Information Science and

Engineering, Dayananda Sagar College of Engineering, Bangalore, India

Prof. V. Suma

Department of Technology and Maritime Innovation, University of Southeast, Horten, Norway

Dr. Noureddine Bouhmala

Go Perception Laboratory, Cornell University, Ithaca, NY, USA

Dr. Haoxiang Wang

Rights and permissions

Reprints and Permissions

Copyright information

About this paper

Cite this paper

Sivapriyan, R., Elangovan, D., Lekhana, K.S.N. (2021). Review of Python for Solar Photovoltaic Systems. In: Suma, V., Bouhmala, N., Wang, H. (eds) Evolutionary Computing and Mobile Sustainable Networks. Lecture Notes on Data Engineering and Communications Technologies, vol 53. Springer, Singapore. https://doi.org/10.1007/978-981-15-5258-8_12

.RIS .ENW .BIB

DOI	Published	Publisher Name

https://doi.org/10.1007/978-981-15-5258-8_12	01 August 2020	Springer, Singapore

Print ISBN	Online ISBN	eBook Packages
978-981-15-5257-1	978-981-15-5258-8	Engineering Engineering (R0)

ENGINEERING CHEMISTRY

ABOUT THE BOOK: Engineering Chemistry aims to impact an in-depth knowledge of various aspects of chemistry to engineering students. It has been structured according to syllabus prescribed by Visvesvaraya Technological University. This textbook contains 5 modules covering various disciplines of Engineering Chemistry.

SALIENT FEATURES:

- The book exclusively covers all the modules of the syllabus.
- Provides a perfect link between the fundamental concepts and their relevant applications.
- Simple and clear language, point-wise description in easy steps so that students can grasp the concepts easily.
- A good number of examples for easy understanding.
- Exercise questions for each chapter are included.
- Clear diagrams and solved numerical problems have been included wherever required.
- Lab manual with viva-voce and their answers are included.

G. M. Krishnaiah is presently working as Professor and HOD of Chemistry Sir MVIT, Bengaluru. He obtained his M.Sc. and Ph.D. degrees in Chemistry from Bangalore University. He has taught Engineering Chemistry for the last 28 years and has published 16 research papers in peer-reviewed journals. At present, he is guiding students for their Ph.D. degree. He served as the member of Board of Studies in Basic Sciences, Ph.D. Registration Expert Committee and Chairman of Board of Examiners for VTU, Belagavi. He was also awarded the best Professor in Chemistry by Dewang Mehta National Education Award.

Prashanth G. K, Assistant Professor in the Department of Chemistry, Sir MVIT, Bengaluru obtained his B. Sc. degree with 2nd place and M.Sc. degree with 3rd rank and gold medals from the University of Mysore. He obtained his M.Phil. degree in Chemistry and presently pursuing Ph.D. degree. He has over 13 years of teaching experience. He has published many research papers in reputed peer-reviewed international journals and presented several research articles in national and international conferences.

NEWRAYS

Website: www.newraysindia.com

MRP. ₹ 399/-

9 788194 068716

ENGINEERING CHEMISTRY

G.M. Krishnaiah
Prashanth G.K.

NEWRAYS



ENGINEERING CHEMISTRY

NEWRAYS

G.M. Krishnaiah
Prashanth G.K.

Performance Evaluation of Nature Inspired Algorithms for MPPT in Solar PV Systems with Partial Shading

Publisher: IEEE Cite This PDF

R Subha. ; S Himavathi. **All Authors**

Abstract: Nature inspired algorithms have been found to be powerful in solving complex optimization problems due to the intelligence embedded in them. Some of these algorithms have... **View more**

Metadata

Abstract:
Nature inspired algorithms have been found to be powerful in solving complex optimization problems due to the intelligence embedded in them. Some of these algorithms have also been used in tracking the maximum power point (MPP) in a partially shaded photovoltaic (PV) array. This paper evaluates the capability of five algorithms inspired from nature in tracking maximum power point under partial shading. The performance of the algorithms are compared in terms of their tracking accuracy, convergence time and implementation complexity to find the most suitable method for MPP tracking.

Date of Conference: 21-23 August 2019 **INSPEC Accession Number:** 19455699

Citations

Keywords

Metrics

More Like This

Date Added to IEEE *Xplore*: 16 March 2020

ISBN Information:

Electronic ISBN:978-1-7281-2414-8

Print on Demand(PoD) ISBN:978-1-7281-2415-5

DOI: 10.1109/ICPEDC47771.2019.9036599

Publisher: IEEE

Conference Location: Chennai, India

Contents

I. Introduction

There has been a phenomenal growth in photovoltaic (PV) power generation in the recent years due to increase in power demand and the benefits of solar power. Multiple peaks in the P-V characteristics during partial shading makes maximum power extraction difficult [1] - [2].Due to multiple peaks, the commonly used algorithms fail to extract maximum power and need modifications for global maximum power point tracking(MPPT) [3]–[5].

Authors

Figures

References

Citations

Keywords

Metrics

Performance of Wind Energy Conversion System during Symmetrical fault & Power Quality Improvement of Grid-Connected WECS by UPFC

Publisher: IEEE Cite This PDF

H L Suresh ; M Sharanappa ; CH Venkat Ramesh **All Authors**

Abstract

Abstract: The demand for the power generation from wind is constantly growing. This situation forces the revision of the grid codes requirements, to remain connected during grid fa... **View more**

Metadata

Abstract:

The demand for the power generation from wind is constantly growing. This situation forces the revision of the grid codes requirements, to remain connected during grid faults. Immediately the voltage level will drop below 80% when fault occurs at PCC (Point of Common Coupling) and the rotor speed of IG (Induction Generators) becomes unstable. In this work, UPFC are used under fault condition to improve the LVRT (Low Voltage Ride-Through) of WECS (wind energy conversion system) and damping of rotor speed oscillations of IG. Furthermore, after the fault UPFC act as virtual inductor, leads to increase in terminal voltage of WECS. WECS with DFIG-based system is considered for analysis here. By simulating DFIG-based WECS with UPFC indicates the improvement in LVRT & remains and WTGs continues to operate with grid at certain voltage fluctuations, near grid. Also, indicates voltage improvement at PCC under fault conduction, and voltage is recovered easily to 1pu at PCC.

Published in: 2018 4th International Conference for Convergence in Technology (I2CT)

Date of Conference: 27-28 October 2018

Date Added to IEEE *Xplore*: 09 April 2020

ISBN Information:

Electronic ISBN:978-1-5386-5232-9

Print on Demand(PoD) ISBN:978-1-5386-5233-6

INSPEC Accession Number: 19514765

DOI: 10.1109/I2CT42659.2018.9058230

Publisher: IEEE

Conference Location: Mangalore, India

Contents

I. Introduction

The energy demand in world increases day by day. To meet this energy demand, we should consider conventional & nonconventional sources of energy. The nonconventional sources of energy are become more important to meet demand in recent years. The nonconventional energies are nothing but Renewable Energy sources like Wind, Solar, and Biomass etc. As compared to all energy sources Wind & Solar abundantly available in nature. But availability wind energy more [1], [2].

Sign in to Continue Reading

Published: 2023

Show More

Recent Research Trends in Solar Photovoltaic Systems

Publisher: IEEE

R. Sivapriyan ; D. Elangovan ; B.S. Kiran ; R. Madan **All Authors**

Abstract: In recent years, conventional energy sources are depleting and getting exhausted. In replacement of this, solar Photovoltaic (PV) systems are used because of its free ava... **View more**

Metadata

Abstract:
In recent years, conventional energy sources are depleting and getting exhausted. In replacement of this, solar Photovoltaic (PV) systems are used because of its free availability, renewabiiity and less operating cost. But the constraints such as low efficiency, dependability on weather conditions are overcome by using maximum power point tracking(MPPT) algorithms and efficient converters to produce the maximum power output. MATLAB/Simuiink is used to model and analyze PV array, MPPT algorithms, and converters. In this paper, recent research developments in PV array, various MPPT techniques, many efficient Converters are presented. This paper will give an overall idea about the present research trends in the field of Solar Photovoltaic systems.

Published in: 2020 5th International Conference on Devices, Circuits and Systems (ICDCS)

Date of Conference: 05-06 March 2020 **INSPEC Accession Number:** 19569023

Metrics

More Like This

Date Added to IEEE *Xplore*: 23 April 2020

ISBN Information:

Electronic ISBN:978-1-7281-6368-0
Print on Demand(PoD) ISBN:978-1-7281-6369-7

ISSN Information:

DOI: 10.1109/ICDCS48716.2020.243584

Publisher: IEEE

Conference Location: Coimbatore, India

Contents

I. Introduction

Nowadays, constant decrease of fossil fuels and increasing demands for energy has led to research on the renewable sources. Solar irradiance is freely available and is the most convenient form of energy to generate electricity. The solar PV array is used to extract electrical energy from the sun light. But due to the change in weather conditions [1], the panel output is low. To improve the output of the solar panel, it is required to operate the PV array at maximum power point (MPP). The MPP technique will adjust the duty cycle of the converters which in turn improves the power from the solar array. To operate the panel at MPP, a well suited algorithm is required. So, by using MPPT algorithm, additional converters and batteries in a PV system, reliable power supply can be provided [2]. This regulated output is used in practical applications. This paper examines the recent research trend in the solar PV system components.

Sign in to Continue Reading

Authors

Figures

References

Citations

Keywords

Metrics

IEEE Transactions on Energy Conversion
Published: 2012

Show More

Access provided by:
Sir M Visvesvaraya Inst. of Technology

Sign Out

Design and Simulation of Mems based Piezoelectric Acoustic sensor

Publisher: IEEE

S. Veena ; Deepika ; Vasudha Hegde ; Veda Sandeep Nagaraj ; H. L Suresh ; M.A. Gautham **All Authors**

Abstract

Abstract:A MEMS based acoustic sensor that combines high sensitivity, wide frequency range and low cost batch processed miniaturized silicon components to build self powered syste... **View more**

Metadata

Abstract:
A MEMS based acoustic sensor that combines high sensitivity, wide frequency range and low cost batch processed miniaturized silicon components to build self powered systems is presented in this paper. It also throws light on an effective method to monitor health of a machine which is by using an piezoelectric Mems microphone. The proposed Acoustic sensor consists of a sputtered piezoelectric ZnO layer that transforms the mechanical deflection of a thin-etched-Si diaphragm into a piezoelectric charge. This ZnO layer is sandwiched between bottom Al electrode and top Al electrode. The simulations of the proposed acoustic sensor is carried out for two designs i) The piezoelectric material being placed at the 4 corners of the silicon substrate and ii) The piezoelectric material being placed at centre of the silicon substrate. The thickness of the layers are chosen so as to withstand the dynamic sound pressure of 96-106db and it produces maximum of 8pV/Pa. The simulation is done by Comsol multiphysics and Coventorware.

Published in: 2019 IEEE 4th International Conference on Condition Assessment Techniques in Electrical Systems (CATCON)

Date of Conference: 21-23 November 2019

Date Added to IEEE *Xplore*: 27 April 2020

INSPEC Accession Number: 19573535

DOI: 10.1109/CATCON47128.2019.CN0061

Publisher: IEEE

ISBN Information:

Electronic ISBN:978-1-7281-4331-6
USB ISBN:978-1-7281-4330-9
Print on Demand(PoD) ISBN:978-1-7281-4332-3

Conference Location: Chennai, India

Contents

I. Introduction

Mechanical energy is one of the most abundantly available and reliable energy sources in our daily life, regardless of weather or temperature conditions, in contrast with solar and thermoelectric energies. At micro scale, energy harvesting technology based on the piezoelectric effect have been developed to harvest small scale and irregular sources of mechanical energy present in our living environment.

Sign in to Continue Reading

Authors

Figures

References

Keywords

Metrics

A REVIEW ON GREEN SYNTHESIS OF SILVER NANOPARTICLE, CHARACTERIZATION AND OPTIMIZATION PARAMETERS

Article · September 2016

DOI: 10.15623/ijret.2016.0527010

Archna Archna
M.S. Ramaiah Institute of Technology

Halima R.
Sir M Visvesvaraya Institute of Technology

A REVIEW ON GREEN SYNTHESIS OF SILVER NANOPARTICLE, CHARACTERIZATION AND OPTIMIZATION PARAMETERS

Halima R[1], Archna[2]

[1]*Department of Biotechnology, Sir M Visvaraya Institute of Technology, Bangalore-562157*
[2]*Department of Chemical Engineering, M.S. Ramaiah Institute of Technology, Bangalore- 560054*

Abstract

Nanotechnology deals with the study of particles in the range of 1-100nm where they have unique properties which define their application. In recent years, green synthesis of silver nanoparticles is prominent with scientists and researches for its ease of availability, nonhazardous waste production and is economically viable too. It is seen that the green synthesis has over taken the chemical synthesis of nanoparticles as it is quick and easy. The methodology of synthesis along with various conditions like the temperature, time, pH, silver ion concentration, radiations also plays a vital role in the shape, size and yield of the nanoparticles produced. Optimization of the parameters for the better quality yield of silver nanoparticles enables us to apply in areas as microbiology, photography, catalysis, biological labeling, phototonics, optoelectronics, medical devices and healthcare products

Keywords: *Nanotechnology; silvernanoparticles; green synthesis; optimization*

--***--

1. INTRODUCTION

In current scenario with a concern to protect the environment, green technology has got its own potential to overcome it. Metal nanoparticles for its own advantage of being used in various applications, the more economic and environment friendly way of producing it is a challenge. The applications of metal nanoparticles are in various field as in electronics as sensors, as an antimicrobial agent, as bio catalytic activity in various reactions, environmental remediation, physics, material science etc. Green synthesis has paved a way of more environmental and economic way of producing silver nanoparticles and with availability of lots of sources it has got its own specificity and application.

Characterization, play a vital role in estimating their nature of application in which the physical properties of the nanoparticles as size, shape, functional groups attached along with the optical properties are known.

The different parameters which influence the synthesis of nanoparticles are the pH, temperature, Silver ion concentration, concentration of the leaf extract used, time of reaction, radiations, pressure and so on. On optimizing these parameters maximum yield with desirable properties can be achieved. This paper discusses on the green synthesis methods of silver nanoparticles with all the characterization methods and the different optimization parameters.

2. GREEN SYNTHESIS METHODS

2.1 Polysaccharide Method

Reduction of silverions to nanoparticle is enabled by polysaccharide as reducing agent that can act as capping agent too. Synthesis of nanoparticle with Alpha D glucose and Beta D glucose in the presence of starch as a capping agent are also carried out. Significance of starch mediated silvernanoparticles is its easy integration into systems of biological and pharmaceutical applications. The interaction between starch and nanoparticles are week and are reversible at elevated temperatures enabling the separation of them (Ravendran et al 2003)

2.2 Tollens Method

The Tollens Synthesis method gives silver nanoparticles with controlled size in about one step process. The basic Tollens reaction involves the reduction of Tollens reagent ($Ag(NH_3)_2$) by an aldehyde. In the modified Tollens procedure Ag^+ ions are reduced by saccharides in the presence of ammonia yield a stable particle size where the ammonia concentration and the nature of the reductant play a major role in controlling the size of silver nanoparicle (Yu etal 2004)

2.3 Irradiation Methodology

Various studies indicate the successful production of silver nanoparticles by radiations.Silvernanoparticles of well defined shape and size can be produced by passing laser irradiation through an aqueous solution of Silver salt and surfactant. Also the microwave radiation of solution containing carboxymethyl cellulose sodium and silver nitrate produced uniform silvernanoparticles stable for 2 months. Stable nanoparticle is also synthesized by irradiating with gamma radiation and passing 6 MeV electrons beam. [26]

2.4 Polyoxometalates Method (POMs)

POMs have the capability of undergoing stepwise, multielectron redox reactions without disturbing their structure. POMs serve both as a photocatalyst, reducing agent and a stabilizer (Bogle etal 2004)

2.5 Biological Method

Extracts from microorganisms and plant, act both as reducing and capping agents in silver nanoparticle synthesis. The reduction process of silver ions takes place by biomolecules such as enzymes, proteins, aminoacid, polysaccharides and vitamins. Lot of researches are going on by using the extracts of various bacterias, fungi, Actinomycetes and also the plant extracts as this seem to be the safest and economical way of producing nanoparticle. Out of this green synthesis, using plant material as source is advantageous overall as they are abundantly available and easy to handle. Different plant sources that produce silver nanoparticles as listed out in Table1.

Table 1 Plant sources for the synthesis of silver nanoparticles with shape and sizes.

Plant source	Part	Size	Shape	Reference
Cinnanmon zeylanicum	Bark	31-40nm	Quasi spherical shaped and rod shaped	(Sathikumar 2009)
Pelargonium graveolens	Leaves	16-40nm	Quasi linear superstructures	(Shanker et al 2003)
Cinnammum camphora	Leaves	55-80nm	Triangular or spherical	(Huang et al 2007)
Aloe vera	Leaves	50nm	Triangular	(Chandran et al 2006)
Cycas	Leaves	2-6 nm	Spherical	(Philip et al 2001)
Jatropha curcas	Latex	20 nm	Spherical	(Bar et al 2009)
Jatropha curcas	Seed	15 to 50 nm	Rod shaped	(Bar et al 2009)
Cynnamon zeylanicum	Bark	80 nm	Spherical	(Sathikumar 2009)
Black Tea	Leaves	40nm	Spherical	(Begum et al 2009)
Desmodium triflorum	Leaves	5-20nm	Spherical	(Ahmed etal 2010)
Murraya koenigii	Leaves	20nm	Hexagonal and spherical	(Jha et al 2009)
Citrus limon	Leaves	50nm	Spherical	(Prathnaa et al 2011)
Coriandrum sativum	Leaves	26nm	Spherical	(Sathyavathi et al 2010)
Capsicum annuum	Leaves	10-12nm	Triangular	(Li et al 2007)
Medicago sativa	Leaves	2-20nm	Spherical	(Gardea et al 2003)
Azadirachta indica	Leaves	5-35nm	Spherical, plate like, core shell structure	(Shankar et al 2004)
Pelargoniumgraveolens	Leaves	16-40nm	Triangular	(Shankar et al 2003)

2.6 Mechanism of Synthesis of Silvernnaoparticles through Plant Extracts

Main mechanism in which the silver nanoparticles are synthesized is through the reduction of silver ions in which the phytochemicals play a vital role in it. Phytochemicals such as terpenoids, flavones, ketones , aldehydes, amides and carboxylic acid play a vital role (Gluskar 1999). Mechanism of synthesis of silver nanoparticles involves three main phases. The first phase is called an activation phase in which the reduction of the silverions is carried out and nucleation of reduced metal ions are carried out. The second phase is called as a growth phase in which small adjacent nanoparticles spontaneously coalesce into particles of larger size (Oswald Ripening). During the growth phase the nanoprticles aggregates to form nanotubes, nanoprisms, nanohexadrons and variety other irregularly shaped nanoparticles. The third phase is called as a termination phase in which the final shape of nanoparticles are confined. Also the nanoparticles acquire most energetically favourable conformation. Plant extracts play a vital role to stabilize the metal nanoparticles (Si etal 2007).

2.7 Role of Plant Metabolites in Silver Nanoparticle Synthesis

Various plant metabolites including terpenoids, polyphenols, sugars, alkaloids, phenolic acids and proteins play an important role in reduction of metal ions which yields nanoparticles. FTIR spectroscopy reveals that Terpenoids are often associated with the nanoparticles. Terpenoids have strong antioxidant activity and was found to play an principal role in bio reduction of silver ions (Rai et al 2007). Flavanoids have various functional group which involves in silver nanoparticle formation. It is noted that the tautomeric transformation of flavanoids from the enol form to keto form may release a reactive hydrogen atom that can reduce metal ions to form nanoparticles.(Ahmed etal 2010) Flavanoids are also able to adsorbed on the surface of nascent nanoparticle. This means they are involved in stages of nucleation and further aggregation in addition to bio reduction stage.

3. CHARACTERIZATION

3.1 Reduction of Silverions

The reduction of silver ions is noticed by the colour change from pale green to dark brown colour which is observed within minutes after the addition of 1mM silver nitrate to the leaf extract. The formation of nanoparticles by different salts gives characteristic peak at different absorptions which can be monitored by UV/VIS spectroscopy. Silver nanoparticle formation from the silverions shows a characteristic peak at 450 nm.

3.2 Shape and Size

Various factors govern the size and shape of the nanoparticle. Estimation of the morphological features silvernanoparticle is by Transmission electron microscope (TEM). Determination of size and shape has a greater impact on its optical behaviour and surface plasmon effect. They can be in triangular, spherical or hexagonal in shape.

3.3 Functional Group

Sometimes the functional group gets attached to the nanoparticle formed. The functional groups attached depend on the source of nanoparticles. This might be also due to the improper purification step performed during synthesis of nanoparticles. The functional group attached to the silver nanoparticle has to be found to ensure its relevance to the application. Functional groups attached to the nanoparticle can be well analyzed by FTIR.

3.4 Crystalline Nature

The crystalline nature of silver nanoparticles can be confirmed by X-ray diffraction studies. The spectrum shows distinct peaks indexed to the reflection of the crystalline planes of face centred cube. Understanding the crystalline nature of silver nanoparticles enables its application in optical sensors and image analysis. The optical properties can be well studies by the addition of stabilizer and capping agent as polyvinyalcohol and glycerol during the synthesis of nanoparticles.

3.5 Size Distribution

Dynamic light scattering system (DLS) is generally used for analysis of the size distribution pattern of particles in suspension or solution which gives the hydrodynamic diameter of particles. The average size of particles in a suspension can be well analyzed by DLS.

3.6 Elemental Composition

Energy dispersive X-ray spectroscopy (EDX) determines the elemental composition. It is used to confirm that the nanoparticle suspension contains nothing but silver.

3.7 Stability Analysis

Zeta potential is an essential parameter for the characterization of stability in aqueous nano suspensions. This possibly gives the charge of the nanoparticles formed. The indication of positive or negative nature of nanoparticles enables its use in electronic devices.

3.8 Thermal Stability

Thermal stability of the nanoparticle formed can be monitored by Thermo Gravimetric Analysis (TGA). The change in weight of the subjected materials with respect to temperature range of 0-1000^{o}C can be carried out. The loss in weight might be related to the decomposition of surface coated phytochemicals, adsorbed water molecules and oxidation of silver nanoparticle upon increase in temperature.

3.9 Silver Ion Concentration Determination

Silver ion concentration can be analysed by Atomic Absorption Spectroscopy (AAS), which shows the conversion of Ag^{+} ions to Ag^{0} nanoparticles. The reaction time of complete conversion of Ag^{+} to Ag^{0} nanoparticles can be well noted by this method.

4. FACTORS INFLUENCING METAL NANOPARTICLE SYNTHESIS

4.1 Silverion Concentration

The concentration of silver nitrate solution used also has an effect on Silver nanoparticle formation. Experiments carried out by varying dilutions of silver nitrate solution also have an effect on the shape and size of the nanoparticle. It is seen that the increased dilution as 1:1 with the extract will give a comparatively bigger size of nanoparticle and lesser concentration gives a smaller sized one. The optimum silver ion concentration at most cases is reported as 1mM.

4.2 Effect of pH

pH has a great influence in formation of nanoparicle. pH has the capability in affecting the phytochemicals by a charge change in them. It also affects the ability to bind to each other during the termination phase and thereby affecting the shape, size and yield of the nanoparticles.

It is well noted that the better yield of nanoparticles are obtained at alkaline pH where is acidic pH low yield or no nanoparticle formation in noticed in different studies.

Table 2 Effect of pH on yield of nanoparticle

pH	Yeild %	Reference
<3	Agglomerates, No yield	(Shankar etal 2003)
3-5	less yield	(Sathikumar 2009)
6	52.3	(Jiang et al 2009)
7	69.8	(Prathna et al 2011)
8	88.7	(Huang et al 2007)
9	86.2	(Gluskar et al 1999)
10	80.3	(Jiang et al 2009)

4.3 Effect of Temperature

Temperature elevation can increase the reaction rate and efficiency of synthesis. Most studies are carried out at room temperature as it is the simplest and natural way of synthesizing the nanoparticle. As the temperature increases the time of reaction decreases and 95% conversion into nanoparticle is obtained at 95^0C within 10 minutes. The temperature greatly influence on the shape of the nanoparticle too. Nanoparticles which give spherical shape at 60^0C are giving triangular shape at 90^0C.

Table 3 Effect of temperature on nanoparticle formation

Temp.	Shape	Plant species	Reference
40^0C	Triangular	*M.sativa*	(Shankar et al 2003)
50^0C	Spherical	*Cassia fistula*	(Prathna et al 2011)
60^0C	Spherical	*Coriandrum sativum*	(Li et al 2007)
70^0C	Spherical,	*C. aromaticus*	(Gluskar et al 1999)
80^0C	Spherical	*Nyctanthes arbortristis*	(Prathna et al 2011)
90^0C	Triangular	*Pelargoniumgraveolens*	(Shankar et al 2003)

4.4. Effect of Time on Reaction

The reduction reaction and formation of nanoparticles starts immediately after the addition of silvernitrate with the plant extract which is noted by the colour change from light green to brown colour. But, it is seen that the particle size increases with increase in time and stabilize at a particular time. For instances studies reveal that seconds after the addition of the plant extract the size was 7nm and further increasing the time nucleation process has proceeded and became stable before 30mins were the size was 20nm.

4.5 Effect of Radiation

It is seen that the radiations have an impact on the silver nanoparticle synthesis. Studies are done on different radiation as Sunlight, Microwave, laser and UV radiation. The radiations induce the synthesis of nanoparticles also contributes on the shape. The sunlight is proved to give greater yield in comparison with other radiation. Studies reveal that the laser radiation produces triangular shaped nanoparticle whereas the microwave radiation has the ability to produce spherical shaped particles.

4.6 Effect of Extract Amount

As the amount of extract added increases the formation of silver nanoparticles increases. The reductase present in the leaf extract enables the synthesis of silver nanoparticles by reduction reaction with silver ions. The ideal concentration is in 1:1 ratio with silver ions. Increase in amount of extract results in the production of larger size nanoparticles.

4.7 Effect of Electron Donor

Glucose (56 mM) acts as an electron donor and has positive effects on nanoparticle synthesis. In absence of electron donor, electron reservoirs in reaction mixture for the recovery and revival of cofactors after a short time are finished and reaction declines.

5. CONCLUSION

Green synthesis of nanoparticles paves a way for future for economical synthesis without any harmful byproduct. Although there are different methods of green synthesis, the biological method using the plant extract serve as the widely used amongst all. It is seen that the source plays a vital role in contributing the shape and size of nanoparticles. As plants serves as a source for nanoparticle production there might be possible contaminations associated with the nanomaterials. Proper purification steps helps in overcoming the contamination. The synthesis of nanomaterials are associated with different factors affecting it such as pH, temperature, reaction time, radiations, silver ion concentration and plant extract. Each factor contributes to the size and shape. The yield can also be increased by optimising certain parameters. Hence it is essential to optimize the synthesis of silver nanoparticles for the better application of the same.

REFERENCES

[1] Ahmad N., Sharma S., Alam M.K., Singh V.N., Shamsi S.F., Mehta B.R., Fatma A.. Colloids Surf. B. Biointerfaces. 2010;81:81–86.

[2] Bar, H., Bhui, D.K., Sahoo, G.P., Sarkar, P., De, S.P. and Misra, A. 2009a. Green synthesis of silver nanoparticles using latex of Jatropha curcas. Colloids Surf A: Physicochem Eng Asp, 339: 134–139.

[3] Bar, H., Bhui, D.K., Sahoo, G.P., Sarkar, P., Pyne, S. and Misra, A. 2009b. Green synthesis of silver nanoparticles using seed extract of Jatropha curcas. Colloids Surf A: Physicochem Eng Asp, 348: 212–216.

[4] Chandran, P.S., Chaudhary, M., Pasricha, R., Ahmad, A. and Sastry, M. 2006. Synthesis of gold nanotriangles and silver nanoparticles using Aloe vera plant extract, Biotechnology Prog., 22: 577-583.

[5] Das R.K., Gogoi N., Bora U.. Bioprocess. Biosyst. Eng. 2011;34(5):615–619.

[6] Gardea-Torresdey, J.L., Gomez, E., Peralta-Videa, J.R., Parsons, J.G., Troiani, H. and Jose-Yacaman, M. 2003. Alfalfa sprouts: a natural source for the synthesis of silver nanoparticles. Langmuir 19:1357–1361.

[7] Glusker J., Katz A., Bock C., Rigaku J. 1999;16(2):8–16.

[8] Huang, J., Li, Q., Sun, D., Lu, Y., Su, Y., Yang, X., Wang, H., Wang, Y., Shao, W., He, N., Hong, J. and Chen, C. 2007. Biosynthesis of silver and gold nanoparticles by novel sundried Cinnamomum camphora leaf. Nanotechnology, 18: 105104-105114
[9] Li, S., Shen, Y., Xie, A., Yu, X., Qiu, L. and Zhang L. 2007. Green synthesis of silver nanoparticles using Capsicum annuum L. extract. Green Chem., 9: 852–885.
[10] Lin L., Wang W., Huang J., Li Q., Sun D., Yang X., Wang H., He N., Wang Y., Chem. Eng. J.2010;162:852–858.
[11] Lukman A.I., Gong B., Marjo C.E., Roessner U., Harris A.T.. J. Colloid. Interface. Sci. 2010;353:433–444.
[12] Prathnaa, T.C., Chandrasekarana, N., Ashok, Raichurb, M. and Amitava Mukherjee. 2011. Biomimetic synthesis of silver nanoparticles by Citrus limon (lemon) aqueous extract and theoretical prediction of particle size. Colloids and Surfaces B: Biointerfaces, 82: 152–159.
[13] Rai M., Yadav A.. IET Nanobiotechnol. 2013;7(3):117–124.
[14] Raveendran P, Fu J, Wallen SL.J Am chem. Soc 2003;125:13940
[15] Sathishkumar, M., Sneha, K., Won, W.S., Cho, C-W., Kim, S. and Yun, Y-S. 2009. Cynamon zeylanicum bark extract and powder mediated green synthesis of nanocrystalline silver particles and its bactericidal activity, Colloids Surfaces, B:Biointerfaces, 73: 332-338.
[16] Shankar, S., Ahmad, A. and Sastry, M. 2003. Geranium leaf assisted biosynthesis of silver nanoparticles, Biotechno. Prog., 19: 1627-1631.
[17] Shankar, S., Rai, A., Ahmad, A. and Sastry, M. 2004. Rapid synthesis of Au, Ag and bimetallic Au core-Ag shell nanoparticles using Neem (Azadirachta indica) leaf broth. J. Colloid Inter. Sci., 275: 496-502.
[18] Si S., Mandal T.K.. Chemistry. 2007;13(11):3160–3168.
[19] Weinstock IA, Chem Rev 1998; 98:113.
[20] Yu D, Yam VW-W J Am Chem Soc 2004; 126:13200.
[21] Begum, A.N., Mondal, S., Basu, S., Laskar, A.R. and Mandal, D. 2009. Biogenic synthesis of Au and Ag nanoparticles using aqueous solutionsof Black Tea leaf extracts, *Colloids Surfaces B. Biointerfaces*, 71: 113-118.
[22] Jha, K.A. and Prasad, K. 2009. Biosynthesis of silver nanoparticles using *Cycas* leaf broth. *Bioprocess Biosystems Eng.*,
[23] Philip, D. 2011b. *Mangifera Indica* leaf-assisted biosynthesis of well-dispersed silver nanoparticles. *Spectrochimica Acta Part A*, 78: 327–331.
[24] Sathyavathi, R., Balamurali Krishna, M., Venugopal Rao, S., Saritha, R. and Narayana Rao, D. 2010. Biosynthesis of Silver Nanoparticles
[25] Using *Coriandrum Sativum* Leaf Extract and Their Application in Nonlinear Optics. *Advanced Science Letters* 3: 1–6.
[26] Zhang, Y., Yang, D., Kong, Y., Wang, X., Pandoli, O. and Gao, G. 2010. Synergetic Antibacterial Effects of Silver Nanoparticles@Aloe Vera Prepared via a Green Method. *Nano Biomed. Eng.*, 2(4): 252-257.
[27] Bogle KA, Dhole SD, Bhorasker VN Nanotechnol2006, 17:3204.
[28] Jiang, J.; Oberdorster, G.; Biswas, P. Characterization of size, surface charge, and agglomeration state of nanoparticle dispersions for toxicological studies. J. Nanopart. Res. 2009, 11, 77- 89.